

LOWER SAN DIEGO RIVER WATER QUALITY 2005 - 2020

Annual (WY20) Water Quality Monitoring Report



West Hills Parkway (WQM Site 11) overpass viewing upstream from USGS gauging station 11022480.

Water Quality Monitoring Results (October 2004 - September 2020)

John C. Kennedy, PE

November 2020

Lower San Diego River Water Quality 2005 - 2020

Table of Contents

Section 1. Introduction	pg 2
Table 1.1 - LSDR Water Quality Index	
Figure 1.1 - Lower SDR Watershed and WQM Sites	
Section 2. Spatial Analysis of WY20 WQM Data and 16-Yr Norms	pg 4
Table 2.1 - Average Annual WQ Data by Individual Monitoring Site	
Table 2.2 - WQ Results by River Reach and Section	
Chart 2.1 - WQ Data Profiles by Site and Reach for This Year and 16-Yr Norms	
Chart 2.2 - WQI Profiles by Site and Reach for This Year and the 16-Yr Norms	
Section 3. Temporal Analysis of WY20 WQM Data and 16-Yr Norms	pg 8
Table 3.1 - WQ Data by Month and Season	
Chart 3.1 - WQ Data Results by Month and Season for This Year and the 15-Yr Norms	
Chart 3.2 - WQI Values by Month and Season for This Year and the 15-Yr Norms	
Section 4. Variance in WQM Metrics (WY05-WY20)	pg 11
Table 4.1 - Running Average WQM Metrics (WY05-WY20)	
Chart 4.1 - Monthly Variance in Temperature and Trends	
Chart 4.2 - Monthly Variance in Specific Conductivity and Trends	
Chart 4.3 - Monthly Variance in pH and Trends	
Chart 4.4 - Monthly Variance in DO and Trends	
Chart 4.5 - Monthly Variance in WQI and Trends	
Chart 4.6 - Monthly Variance in Rainfall, Streamflow and Trends	
Section 5. Trends in LSDR WQI (WY05-WY20)	pg 16
Table 5.1 - Average Annual and Seasonal WQI by Reach and Section	
Chart 5.1 - Upper Santee Basin WQI Trends (Oct. 2004 - Sept. 2020)	
Chart 5.2 - Lower Santee Basin WQI Trends (Oct. 2004 - Sept. 2020)	
Chart 5.3 - Mission Gorge WQI Trends (Oct. 2004 - Sept. 2020)	
Chart 5.4 - Upper Mission Valley WQI Trends (Oct. 2004 - Sept. 2020)	
Chart 5.5 - Lower Mission Valley WQI Trends (Oct. 2004 - Sept. 2020)	
Chart 5.6 - Lower San Diego River Watershed WQI Trend Lines (Oct. 2004 - Sept. 2020)	
Appendices: ^(a)	
A. RiverWatch WQM Program Volunteers	A-1
B. Glossary	B-1
C. References	C1-C2
D. LSDR WQM Data Summary (WY05-WY20)	D1-D2
E. LSDR RiverWatch WQM Program	E1-E4
F. LSDR Stream Flow and Water Quality	F1-F3
G. WY19 Monthly WQM Data by Monitoring Site	G1-G9
H. Water Quality Indexing and 2020 WQIs by Monitoring Site (SDRPF)	H1-H5

^(a) Appendices A-H are contained in a separate document.

Questions regarding the San Diego RiverWatch WQM database or interpretation of results expressed in this and similar SDR WQ data monitoring reports can be directed to the attention of John C. Kennedy, through contacting SDRPF at info@SanDiegoRiver.org or the WQM Coordinator, Shannon-Quigley Raymond, at 619-297-7380.

Section 1 - Introduction

This report provides a summary of monthly values, seasonal patterns and annual trends in water quality monitoring data gathered and evaluated by SDRPF's RiverWatch citizen volunteers. WQM data collected monthly over the past 16 years at 15 monitoring sites within the Lower San Diego River (LSDR) watershed have been aggregated, in conjunction with hydrologic streamflow data to develop a numeric water quality index (WQI). Basic monthly data regarding individual water quality parameters and river hydrology for each of the sites monitored are maintained in an extensive Excel database file available at the SDRPF offices; this annual report examines Water Year 2020 (WY20) data in comparison to previous year results and 16-yr averages (norms). The LSDR water quality monitoring site locations are shown on **Figure 1-1**.



Figure 1-1 LSDR Watershed and Water Quality Monitoring Sites

Color Code for LSDR reaches on Figure 1-1 above: Estuary (orange), Lower Mission Valley (purple), Upper Mission Valley (red), Mission Gorge (green), Lower Santee Basin (pink), Upper Santee Basin (dark blue), Lakeside to El Capitan Reservoir (light green) and principal tributaries (light blue)

The water quality sites on Figure 1-1 and monthly RiverWatch water quality data can be viewed in detail on the SDRPF RiverWatch Online Information Center webpage available at <www.sandiegoriver/river_watch.html>. Clicking on the right hand side of the page allows access to the data portal. In addition to water quality monitoring data, the portal also contains: San Diego StreamTeam Bio-assessment data, 401 Project information and USGS real-time streamflow data regarding daily peak discharge and gauge height for the two San Diego River gauging stations (Fashion Valley & Mast Rd Bridge near Santee).

The SDRPF water quality index (WQI) represents the RiverWatch team’s response to the public’s general questions and concerns regarding overall health of the Lower San Diego River system. The index is a numeric (0-100) whereby increasing values signify improving water quality. The numerical index incorporates basic physical, chemical and bacteriological water quality data by integrating six parameters: water temperature (Temp), pH, specific conductivity (SpC), dissolved oxygen (DO and/or %DOSat), mean coliform count (MCC) and streamflow (Q); through determination of weighted factors for each parameter. The resulting values are aggregated to arrive at an overall score for each site, reach, section as well as the entire lower watershed (LSDR). The index values, grade, color codes and general conventions employed are presented in **Table 1.1**.

Table 1.1 LSDR Water Quality Index

SDR WQI (0-100)	Grade	Color Code	Percentile Range	Water Quality Threshold	General
75 or >	A - Very Good	Dark Blue	25%	Well above acceptable WQ criteria	Healthy (>50)
50 - 74	B - Good	Light Blue	25%	Exceeds acceptable WQ criteria	
38 - 49	C - Fair	Green	12.5%	Meets many but not all WQ criteria	Impaired/Ailing (25-49)
25 - 37	D - Marginal	Yellow	12.5%	Meets some acceptable WQ criteria	
13 - 24	E - Poor	Brown	12.5%	Meets few minimum WQ criteria	Unhealthy (< 25)
0 - 12	F - Very Poor	Pink/Rose	12.5%	Well below minimum WQ criteria	

Note: The WQI was developed for fresh water quality metrics only and is not applicable for estuarine or ocean waters.

In general, sites with WQI values of 50 or above (blue zone) exceed expectations for acceptable water quality and are indicative of relatively ‘healthy’ conditions. Scores between 25 and 49 (yellow zone) describe ‘impaired or ailing’ quality levels where evidence exists regarding failure to meet minimum water quality criteria. Waters’ with scores of less than 25 (red zone) do not meet minimum expectations and are considered ‘unhealthy’ and highly stressful to many aquatic life forms. For WQ parameters monitored by RiverWatch, the index expresses results relative to those levels necessary to sustain designated beneficial water uses for the LSDR (Hydrologic Area 907.1) based on State of California Water Quality Standards. Where criteria are non-specific, results are expressed relative to Southern California coastal area freshwater objectives. The inland freshwater index does not apply to tidal estuaries or ocean waters.

Index values were computed using two formulas; one involving four parameters (Temp, SpC and DO) monitored by RiverWatch combined with streamflow (Q); the second with two additional parameters (pH and MCC) combined with averaged streamflow. The equations used for both formulas (WQI₄ and WQI₆) are presented in Appendix B. Differences between the two determinations were found to be minor. The initial determination (WQI₄) presents a broader range of values than the second, as the ‘normalizing’ effects of pH and MCC values (both of which present less spatial and temporal variances for the LSDR) are excluded. The broader range WQI₄ values are expressed in this and previous annual reports.

The index, although specifically developed for the Lower San Diego River, might also be applied to other California coastal watercourses where comparable water quality metrics (i.e., DO, SpC, water temperature and streamflow) are monitored on a regular and consistent basis. A report comparing relative water qualities in three San Diego County watercourses; Los Penasquitos Creek below Poway, the Santa Margarita River below Temecula and near Fallbrook (SUMP), and the Lower San Diego River near Santee and in Mission Valley, prepared through the SDRPF RiverWatch program, is on file.

Section 2 - Spatial Analysis of WY20 Water Quality Metrics and 16-yr Norms

Monthly water quality data collected and recorded at each site by RiverWatch WQM Team volunteers are used to determine annual averages, seasonal patterns and trends as presented in this annual report (and appendices). Supporting streamflow data collected by USGS are also included in the computations. The annual average water quality values for each of the monitoring sites for WY20 as well as the 16-yr norms (average values over the past 16 years of monitoring) are presented in **Table 2.1**. WY20 values greater than the norms are expressed in blue, whereas WY20 values below norms are expressed in red.

Table 2.1 Average Annual WQ Metrics for WY20 and 16-yr Norms by Site, Reach and Section

Site:	LSDR Reach & Section		Temp, oC	SpC, mS/cm	pH	DO, mg/L	DO %Sat	Flow, cfs	WQI ⁴ Values ^a , (Difference) & Grade	
1	LMV	West	20.7/19.7	2.6/2.7	7.8/7.8	6.5/6.2	72/67	49/30	38/37 (+1)	C-/D+
2			19.5/19.0	2.5/2.6	7.8/7.7	5.0/4.4	54/46		34/30 (+4)	D/D
3			19.7/19.2	2.5/2.5	7.9/7.8	5.3/4.6	57/49		36/31 (+5)	D/D
4			19.7/19.7	2.2/2.5	7.9/7.8	6.2/6.1	68/66		41/40 (+1)	C/C
5	UMV	West	17.9/17.2	2.5/2.6	7.7/7.6	4.8/4.8	50/49	46/28	33/32 (+1)	D/D
6			18.8/18.3	2.4/2.6	7.6/7.6	3.6/3.6	36/36		26/25 (+1)	D-/D-
7			18.9/18.1	2.3/2.5	7.7/7.6	6.3/5.1	66/52		42/34 (+8)	C/D
8	MG	Mid	17.8/17.2	2.2/2.3	8.0/7.7	7.5/7.3	79/74	26/19	51/47 (+4)	B-/C
9 ^b			16.2/15.8	4.0/4.9	8.4/7.9	9.7/9.2	98/94		38/36 (+2)	C-/D
10			17.4/17.7	2.1/2.3	7.8/7.8	6.2/7.0	63/73	21/16	37/44 (-7)	D+/C
11	LSB	East	16.9/16.7	2.2/2.2	7.9/7.9	6.1/6.1	63/59			41/37 (+4)
12 ^b			17.7/17.7	1.2/1.6	8.4/7.9	7.2/7.1	76/72		43/36 (+7)	C/D
15 ^b			17.3/18.0	2.3/2.7	8.1/8.1	7.0/7.0	73/71	15/10	42/39 (+3)	C/C
13	USB	East	17.6/18.4	1.9/1.9	7.6/7.7	1.2/2.8	14/29	9/5	8/16 (-8)	F/E-
14			19.4/17.6	1.4/1.5	8.0/7.8	4.5/3.4	49/34		28/19 (+9)	D/E
(1-15)	LSDR Avg.		18.3/18.0	2.2/2.3	7.8/7.7	5.4/5.3	56/54	33/23	34/33 (+1)	D/D
c	LSDR (Qwt)		18.4/18.0	2.1/2.3	7.8/7.7	5.0/5.0	53/51	30/20	32/31 (+1)	D/D

a) Average annual water quality index values, change (+/-) and resultant WQ letter grade for WY20 (bold) and the 16-yr norms (italics); values below norms for each metric are in red; values above norms in blue.

b) Lower San Diego River water quality monitoring sites located on tributary (T) streams.

c) Average flow-weighted LSDR WQ Index values based on USGS streamflow data presented in Appendix H.

All but two (10&13) monitoring sites present WY20 average annual WQI values greater than their 16-yr norms. Average annual water temperatures in WY20 exceeded 16-yr norms at 12 out of 15 sites; while overall up 0.3 degree from the 16-yr annual average of 18.0 C. Specific Conductivity values in WY20 were equal (3,11&13) or slightly below 16-yr norms at all monitoring sites. Overall SpC (LSDR average) is only 4% less the 16-yr average annual norm of 2.29 mS/cm. DO values are greater than 16-yr norms at 8 sites, same at four and less at two (10&13). Overall this year’s average DO values are approximately 1% higher than the 16-yr LSDR annual norm (5.3 mg/L/54%Sat). DO values for WY20 are also above those from last year (WY19) by nearly 0.5 mg/L (3%Sat). They are well above the poorest year (WY14) by nearly 1.5 mg/L. The highest average annual DO levels were monitored in WY05 at 6.6 mg/L (62% Sat.).

Average annual, seasonal and monthly min.-max. range water quality metrics for WY20 and the 16-yr norms are also presented by river reach and section in **Table 2.2**. Two reaches of the river (USB & MG) present slightly lower index values for the past year than associated 16-yr norms. Average annual water temperatures, dissolved oxygen and specific conductivities for all reaches and sections of the river were very near 16-yr norms. Streamflow exceeded 16-yr norms in all reaches and sections throughout WY20. The most significant improvements in water quality metrics monitored within the lower river watershed occurred during the dry-weather months in the Lower Santee Basin (sites 11,12&15) and Upper Mission Valley (sites 5,6&7) reaches. The least improvement in water quality was monitored in the Upper Santee Basin (sites 13&14).

Table 2.2 Water Quality Metrics for WY20 and 16-yr Norms by Season, Reach and Section



Parameter, units	Temp, oC	SpC, mS/cm	pH	DO, mg/L	DO %Sat	Flow, cfs	WQI Value, ^a (Diff) & Grade		
Max. Month	25.2/25.4	3.1/4.0	8.6/8.7	7.8/10.2	85/103	167/230	52/78 (-26)	B-/A-	
Winter (D,J,F,M)	13.6/13.6	1.3/1.7	7.7/7.7	7.2/6.9	70/66	18/45	49/47 (+2)	C+/C	
Avg. Annual	18.3/18.0	2.2/2.3	7.7/7.7	5.4/5.3	56/54	33/23	34/33 (+1)	D/D	
Avg. (Qwtd)	18.4/18.0	2.2/2.3	7.7/7.7	5.0/5.0	53/51	30/20	32/31 (+1)	D/D	
Summer (J,J,A,S)	23.3/22.5	2.7/2.8	7.7/7.7	4.2/3.8	49/43	2.7/2.1	22/19 (+3)	E/E	
Min. Month	10.9/9.3	0.6/0.6	7.5/7.1	3.6/1.9	38/23	0.9/0.1	24/7 (+17)	E+/F	
<i>LSDR Reach & Section Averages:</i>									
USB	East	18.2/18.1	1.5/1.8	7.8/7.7	2.3/3.0	25/31	9/5	15/17 (-2)	E/E
LSB		17.5/17.4	2.1/2.3	7.8/7.8	6.5/6.5	67/64	21/16	41/37 (+4)	C/D+
MG	Mid	17.3/17.1	2.1/2.3	8.1/7.8	7.5/7.5	77/77	26/19	44/46 (-2)	C/C
UMV	West	18.5/17.9	2.5/2.5	7.7/7.6	4.9/4.5	51/46	46/28	33/30 (+3)	D/D
LMV		19.9/19.4	2.6/2.6	7.8/7.8	5.5/5.1	60/54	49/30	37/35 (+2)	D+/D

a) Average annual water quality index value, difference (+/-) from 16-year norms and resultant WQI letter grades. Values and grades below 16-year norms (shown in italics) are expressed in red; values above in blue.

Spatial water quality values expressed in Tables 2.1 and 2.2 for the fifteen Lower San Diego River system monitoring sites are presented in **Chart 2.1** (Water Quality Data Profiles) and **Chart 2.2** (Water Quality Index and LSDR Streamflow) on the following page. The overall water quality index for WY20 of 32 (D Marginal) is one point above the 16-yr average annual norm of 31 (D Marginal). This year's average annual index value is 10 points above the lowest annual WQI of 22 (E Poor) experienced in WY14. The river's highest overall average annual index of 41 (Fair) occurred in WY05. Only two water year's (WY14 and WY18) have shown overall average index values in the Poor E (WQI 13-24) range.

Average annual water quality values for water temperature, pH, dissolved oxygen and specific conductivity at each monitoring site, river reach and section in order of their location upstream for WY20 (Oct.'19-Sept.'20) and the 16-yr norms are expressed in **Chart 2.1**. This year's average annual results are shown as heavy solid lines with values presented; blue lines are last year's (WY19) results and the red lines are 16-yr annual averages (or norms) for each site. Average annual water temperatures (solid red line) for WY20 are slightly above both 16-yr norms (red bars) and last year values (dashed red line) at nearly all monitoring sites. Average water temperatures downstream are generally higher than monitored at upstream sites. There is little variance in average pH values between sites, from one year to the next or from the 16-yr norms (yellow bars). DO levels for WY20 (solid black line) are generally above those from last year (dashed black line) and near the 16-yr norms (blue bars). Average annual DO values at two sites (6&13) were below threshold levels of 4 mg/L; whereas five sites showed averages below 4 mg/L in WY18. Monitored DO values represent the greatest variation between sites. Lowest values are typically recorded in the Upper Santee Basin (sites 13&14) and Upper Mission Valley (sites 5& 6) whereas the highest values are observed in the Mission Gorge section (middle reach, sites 8-10). Excluding tributary sites, average annual conductivity (SpC) values generally increase along the mainstem from upstream to downstream. SpC averages for WY20 (solid blue line) are near 16-yr norms (brown bars) and last year's values (dashed blue line) at all sites. The greatest variances in this year's spacial metrics both from last year (WY19) and the 16-yr norms are associated with dissolved oxygen and water temperature values.

The WQI, an aggregate or composite index of water quality monitoring metrics for WY20, the 16-yr norms, the overall best (WY05) and worst (WY14) year results are presented in **Chart 2.2**. As shown by the solid black line (this year's results) in comparison to the colored bars (16-yr norms), the two sites furthest upstream, Mast Park (13) and Magnolia Ave (14), continue to experience Poor (E) to Very Poor (F) water quality as does the Kaiser Ponds (site 6). On an average annual basis, highest WQI values continue to be associated with the three Mission Gorge sites (8-10). The overall WQI profile for WY20 (black line) is in general near the 16-yr norms (colored bars) and slightly above last year's (WY19) results (dashed black line). Greatest departures (variance) from the 16-yr WQI norms for WY20 are found at OMD (site 10) and Mast Park East (site 13). Water quality conditions throughout Mission Valley (both Upper and Lower reaches) in WY20 are noticeably improved from last year's (WY19) values. As evidenced in the past, above normal flow tends to flush (or purge) the lower river system resulting in improved overall water quality. WY20 experienced above normal streamflow overall resulting in enhanced base flows throughout the dry-weather months that were instrumental in maintaining above average water quality index values for the entire water year at nearly all monitoring sites with exception of Mast Park East (site 13).



Section 3 - Temporal Analysis of WY20 Data and 16-yr Norms

Monthly, seasonal and annual water quality monitoring metrics data and WQIndex results for the Lower San Diego River are presented in **Table 3.1** for this year (WY20) with comparison to 16-yr norms (shown italicized). WY20 values above the 16-yr norms are in blue; values below in red. With few exceptions temporal water quality WY20 values exceeded 16yr norms on an annual basis for water temperatures, pH DO, streamflow and WQI values while annual SpC were lower. Individual months and seasonal averages varried above (blue) and below (red) 16-yr norms significantly.



Table 3.1 Average LSDR WQM Metrics for WY20 and 16-yr Norms by Month and Season

		Temp	SpC	pH	DO	DO%	Flow	WQI ^(a)	
Month	Season:	<i>oC</i>	<i>mS/cm</i>		<i>mg/L</i>	<i>%Sat</i>	<i>cfs</i>	<i>Value & Grade</i>	
Oct	Fall	18.7/18.7	3.05/2.85	7.6/7.7	3.36 /3.67	37/39	1.0/2.1	16 /20	E/E
Nov		16.1/14.9	3.12 /2.72	7.5/7.7	3.45 /4.92	35/48	7.9 /8.0	20 /28	E/D
Dec	Winter	10.9 /11.9	1.80 /1.80	7.8/7.7	7.45 /6.20	70/57	62/28	47 /41	C/C
Jan		11.9/11.9	1.72/1.70	7.8/7.7	7.28 /7.50	68/69	46/53	50 /48	B-/C+
Feb		14.7/14.0	1.93/1.64	7.8/7.8	6.15/6.84	61/66	9.4 /57	43 /45	C/C
Mar		16.8/16.7	1.10 /1.57	7.9/7.8	6.25 /6.50	64/66	39 /49	48 /49	C+/C+
Apr	Spring	17.4 /18.0	0.88 /1.87	7.8/7.7	5.77/5.40	60/57	167/25	45 /40	C/C
May		19.7 /20.0	1.70 /2.17	7.9/7.7	4.60 /4.69	51/52	18/11	35/33	D/D
June	Summer	21.8 /22.0	2.18 /2.52	7.8/7.7	3.63 /3.96	42/45	6.2/3.9	23 /24	E+/E+
July		24.3/23.2	2.61 /2.72	7.9/7.7	3.64 /3.16	44/37	2.5/2.0	20 /17	E/E
Aug		25.2 /23.4	2.87 /2.94	7.8/7.7	4.44 /3.08	55/36	1.1 /1.2	20 /15	E/E-
Sept		22.0/21.4	2.80 /2.96	8.1/7.7	4.08 /3.08	43/35	0.9 /1.3	19 /16	E/E
Fall (O&N)		17.5/16.8	3.08 /2.78	7.6/7.7	3.66 /4.31	38/44	1.3 /5.0	19 /24	E/E+
Winter (D,J,F,M)		13.6/13.6	1.64 /1.68	7.8/7.7	7.21 /6.79	70/65	63/47	49 /46	C+/C
Spring (A&M)		18.5 /19.0	1.29 /2.02	7.9/7.7	5.59/5.12	60/55	2.0 /18	43 /36	C/D+
Summer (J,J,A,S)		23.5 /22.5	2.61 /2.79	7.9/7.7	4.47 /3.42	53/39	2.4 /2.1	22 /18	E/E
Annual (O-S)		18.4 /18.0	2.15 /2.28	7.8/7.7	5.01 /4.92	53/51	30.1 /20	32 /31	D/D

a) Values based on RiverWatch physical-chemical metrics (WQI₄) combined with USGS stream flow for eastern (West Hills Pkwy) and western sections (Fashion Valley). WY20 values/grades below 16-yr norms (in italics) are shown in red; those equal to or above norms in blue.

Monthly and seasonal variances in water quality monitoring metrics for the past two water years and the 16-yr norms are expressed in **Chart 3.1**. (WQM Data) on the following page. Dissolved oxygen values are highest during the winter/spring months (Dec-May) whereas specific conductivity and water temperatures are greatest during the dry summer months (June-Sept) and into early Fall (Oct). pH values show less seasonal fluctuation, although some variance from norms in the monthly values are evident. The broad range in DO, SpC and temperature values monitored at all sites throughout the year provide the best indications of temporal variance in water quality. Seasonal variances between this year's data (WY20), shown as solid lines, last year's results (dashed lines) and the 16-yr norms (bars) are comparable. In general, temporal variance in WY20 water quality data closely match patterns in 16-yr norms, slightly more so than last year's values. This year's temporal water quality metrics are indicative of both normalized monthly occurrences as well as those monitored during the previous year (WY19). The greatest distinction between last year's metrics and this year's occur during the dry-weather (summer) season. Streamflows, as shown on the next chart, present a significant impact on variances in the other temporal WQ metrics.

Chart 3.2 provides an overall graphic showing temporal variance in WQI values and streamflow throughout WY20 compared to monthly averages over the previous water year (WY19) as well as the 16-year norms. As shown in **Chart 3.2**, the WQI values for WY20 (heavy red line) that are also listed in Table 3.1 (far left column) are relatively close to the 16-yr norms (colored bars) for most months of the year. The strong relationship between flow (both wet weather and dry) and water quality continues to effect results. Somewhat greater DO levels throughout the Winter, Spring and Summer months combined with above normal dry-weather flows constitute the primary drivers in elevated index values. In general, water quality for the Lower San Diego River watershed is highest (B-C Good to Fair) when flows are greatest during the Winter months (Dec-March) and poorest (E-F Poor to Very Poor) in Summer (June-Sept) when streamflow is lowest and water temperatures highest. The overall annual average WQI for the LSDR in WY20 of 32 (D mid-Marginal) is one point above the 16-yr average index value of 31.3. Last year's one point (30) below normal results occurred during a somewhat below average rainfall and runoff year. Although DO depletions were not as great this year as last they have persisted at several sites throughout the dry-weather period, particularly in the upper Santee Basin where streamflow throughout the Summer (June-Sept) remained minimal.



Section 4 - Trends in Water Quality Metrics (WY05 through WY20)

Trends in SDRPF monitored water quality metrics, based on data collected by RiverWatch from September 2005 through September 2019, are presented in this chapter. The metrics include water temperature, specific conductivity, pH, dissolved oxygen, streamflow and the water quality index. Twelve month running average values considered with overall straight-line averages represent a reasonable indication of relative change over the past 15 years of monitoring for each metric.

Table 4.1 presents 12-month running average values for each of the key water quality metrics monitored over the last 16 years. Running averages above 16-yr norms are listed in blue; values below norms are in red. The 16-yr norms (also 12-mo running averages) are expressed in italics in the bottom row.

Table 4.1 - 12-mo Running Average WQM Metrics (WY05-WY20)

	Temp	SpC	pH	DO	DO%	Flow	WQI ^(a)	
	<i>oC</i>	<i>mS/cm</i>		<i>mg/L</i>	<i>%Sat</i>	<i>cfs</i>	<i>Values & Grade</i>	
WY05	17.70	2.064	7.63	6.63	62	71.5	41/40	C Fair
WY06	18.32	2.141	7.44	6.00	59	13.6	36/35	D+ Marginal
WY07	17.79	2.342	7.73	5.95	60	9.5	36/35	D+ Marginal
WY08	17.83	2.223	7.89	6.26	63	18.2	37/37	C- Fair
WY09	17.85	2.393	7.66	6.25	64	20.1	36/36	D+ Marginal
WY10	18.10	2.287	7.84	5.21	54	32.4	34/32	D Marginal
WY11	17.79	2.160	7.83	5.53	57	46.9	38/36	C- Fair
WY12	18.04	2.339	7.64	5.16	53	14.9	33/31	D Marginal
WY13	17.40	2.441	7.77	5.30	54	9.1	32/30	D Marginal
WY14	17.91	2.505	7.67	3.87	40	5.1	22/20	E Poor
WY15	18.60	2.189	7.77	4.53	48	10.t	29/25	D Marginal
WY16	18.19	2.269	7.71	4.69	49	15.6	28/27	D Marginal
WY17	18.57	2.154	7.78	5.05	53	40.0	33/31	D Marginal
WY18	18.31	2.788	7.94	4.28	44	5.9	24/22	E Poor
WY19	17.90	2.170	7.71	4.91	51	26.9	31/30	D Marginal
WY20	18.35	2.154	7.77	5.35	56	32.1	34/32	D Marginal
<i>16-yr Avg</i>	<i>18.04</i>	<i>2.289</i>	<i>7.72</i>	<i>5.31</i>	<i>54</i>	<i>23.3</i>	<i>33/31</i>	<i>(D Marginal)</i>

Values based on SD RiverWatch physical-chemical metrics (WQI₄) combined with USGS streamflow for eastern (West Hills Pkwy) and western (Fashion Valley) gauging stations.




Running average, maximum and minimum monthly monitoring site **water temperature** results for the LSDR watershed are presented on **Chart 4.1**. Running average water temperatures held fairly steady between WY05 and mid-WY11, declined by approximately 0.5 oC by WY14, increased by approximately a degree over the next year (by late WY15) and remained reasonably constant during the last five years. Typical variances in running average water temperature over the past decade are in the range of 3% above to 3% below norms, however, from Oct. 2013 to Oct. 2015 (24 months) variance in water temperature rose from 4.8% below to 5.0% above the 16-yr norm of 18.0°C. Max. monthly water temperatures have trended slightly higher than monthly minimums over the past decade. Higher running average water temperatures observed over the past few years are a result of higher 24-hr average, daytime and nighttime lows in both air and ground temperatures experienced in San Diego as well as throughout much of Southern California. As can be seen in both running average colored lines for max (red) and average (black) and associated dashed lines that trends in water temperature over the past 16 years are slightly upward. The average annual increase is on the order of 0.4%; an overall rise in average annual river temperatures of approximately 0.5°C. Average water temperature for the LSDR in WY20 was 18.4°C, up 0.5 oC from last year's value. The coolest water temperature year since monitoring began was WY13 when the average annual water temperature was 17.4°C. WY15 was the warmest water temperature year at 18.6°C.

Trends in monthly monitored **Specific Conductivity** (SpC) values for the LSDR are presented in **Chart 4.2**. Min. and max. running averages for all sites monitored have varied little over the 16-yr period, however, the overall LSDR running average rose from a low 2.06 mS/cm range (*10% below average*) during the first few years of monitoring to 2.78 mS/cm (*21% above average*) two years ago (WY18). Considerably greater rainfall during WY19 and WY20 and resultant above normal dry-weather stream flow have caused SpC values to fall below the 16-yr norm of 2.28 mS/cm. The current LSDR running average SpC of 2.150 mS/cm is 5.9% below the 16-yr norm. The overall trend in SpC for all sections of the river has shown slight decline over the past 5 years. The variance in maxima at all sites has declined fairly steady of the past 16 years of monitoring, however, site minimum values have also fallen due to less average streamflow (upstream) and rising daily temperatures, resulting in higher evaporation rates.

Trends in monthly **pH** values are presented in **Chart 4.3**. The overall or general trend in values monitored for the LSDR has been relatively consistent over the 16 years (WY05-WY20). The initial five years of below average pH may have been due, at least in part, to faulty equipment as monthly minima and maxima values since WY10 have consistently read higher. Excluding the initial year's, there has been but a small variance (<3%) in the overall running average pH from the 16-yr norm of 7.71. The overall trend in pH for the lower river is, however, slightly positive. Values have increased by an average of 0.3% per annum since RiverWatch monitoring was started, primarily as minima values have risen. It is concluded that the lower river may be becoming slightly more alkaline (basic) as average flows have declined and water temperatures have increased. The most common cause of higher pH water is less available carbon dioxide caused by elevated rates of aerobic respiration that accompany warmer waters. Tracking the trend in pH is important as a general indicator of the natural process of eutrophication in the lower sections of the river.

Running average **dissolved oxygen (DO)** values and monthly minima-maxima are presented in **Chart 4.4** (pg14). A general but somewhat irregular decline in average and min/max values from Oct. 2004 through 2015 can be observed. LSDR maximum monthly values between WY16 and this year have slowly increased to near 16-yr norms. The current running average DO value of 5.01 mg/L (Sept 2020) is 1% (0.03 mg/L) above the 16-yr norm of 4.98 mg/L. Depleted dissolved oxygen levels that have been monitored throughout various reaches and segments of the lower river result from low streamflow, especially during the driest-weather months, combined with above average water temperatures and rapid decomposition of oxygen demanding organic materials (biomass). With a lack of significant flushing action during

Lower San Diego River Water Quality Monitoring Report



relatively mild storm flow events over the past decade, a large amount of decomposing biomass* has accrued within slower moving portions of the river. Overall running average DO values can be expected to improve subsequent to one or more major stormflow events resulting in significant channel flushing, displacement of organic-rich sediments and significant reduction of poorly-rooted and free-floating invasive aquatic plants. The trend in overall LSDR DO values has, over the past 16 years, declined in excess of 2 mg/L from roughly 6.5 mg/L to 5.0 mg/L. This represents an average annual drop in DO of approximately 2.4% (0.13 mg/L) since RiverWatch monitoring was initiated. As can be seen on Chart 4.4, the rate of decline in minimum values (-3 % per annum) is noticeably greater than the rate of decline in maxima (-0.5%/yr). Extended periods of low flow have resulted in lower overall DO levels. Minima values continue to decline at greater rates than maximas.

Trends for total monthly rainfall and running average streamflows in the Santee Basin (SB) and Mission Valley (MV) sections are expressed in **Chart 4.5**. The trend in average daily streamflow throughout the LSDR watershed fell by an order of magnitude (from 57 cfs to 9 cfs) between WY05 to WY07, then slowly rose to 42 cfs by WY11. Lowest running average streamflows of 7-8 cfs for Mission Valley and 3 cfs for the Santee Basin, occurred in WY14. Due to the distribution and magnitude of rainfall in both WY15 and WY16, running average streamflows rose back to 15-20 cfs (Mission Valley) and 8-12 cfs (Santee Basin), but still below 16-yr norms. WY18 streamflows fell sharply as the watershed received near record low rainfall. Dry weather flows in June through September of last year were some of the lowest recorded in the past 4-5 decades. With above normal rainfall in WY19 and WY20, streamflows at both sites climbed back to above long-term norms.

The overall **water quality index** (WQI) for LSDR as well as minimum and maximum running average values for monitoring sites within the watershed are presented in **Chart 4.6**. The WQI provides a general indication of the relative condition of the river based on individual water quality parameters monitored by RiverWatch and streamflow (river discharge) as measured by the USGS at their two gauging stations. Similar to trends in DO (Chart 4.4), running average WQI values that were in general decline from late WY09 to early WY15 slowly increased through 2017. LSDR running averages reached their lowest value of 20 (E Poor) in 2014, at 35% below the 16-yr norm of 31 (D Marginal). WY18 presented the second lowest index at 22, 28% below the norm. This year's running average WQI of 32 (D Marginal) is 3% above the norm. Above normal rainfall (and when it occurred) this year resulted in running average index values similar to those experienced in WY10. A below average rainfall year next year could result in a decline in the index. Much depends on hydrodynamics of the river both during wet and dry-weather periods. A major flushing flow at some point in time could also have a significant impact on the index trend. Over the past 16 years the index has fallen roughly ten points or an average of 0.65 points per annum. Both minima and maxima index values have fallen at comparable rates.

The trends and relative variances in water quality metrics shown in **Charts 4.1-4.6** are interrelated. Less rainfall results in less streamflow (runoff) which results in declining dissolved oxygen levels and increased specific conductivities. As all of the parameters are incorporated in computation of the water quality index, trends over the 16 year monitoring period are similar. The lower river system experienced its best water quality during the wettest year (WY05) followed by a general decline during the well-below average rainfall and river discharge period from WY10 through WY16. The river experienced its poorest water quality during the driest, lowest average annual streamflow year (WY14) recorded over the past 16 years. An uptrend toward normalized values was evident over the past several years (WY15-WY17), but again declined in WY18. WY19 and WY20 have witnessed some recovery. WQI trendlines by individual river reach and specific segment as well as for the overall river system are presented in Section 5.

Section 5 - Trends in LSDR Water Quality Index (WY05 through WY20)

Annual and seasonal LSDR WQI values are presented in **Table 5.1** by river reach, section, and overall (LSDR) average for each water year (WY05-WY20) of monitoring. Values and grades above 16-yr norms are listed in black; values below the 16-yr norms (expressed in italics) are shown in red. The WY20 values, expressed in bold font, are improved over last year's results for all reaches and sections of the lower river. Overall the LSDR average annual WQI rose seven points from last year's value increasing from the E+ Poor water quality range to D Marginal, two points below the 16-year norm.

Table 5.1 - Average Annual and Seasonal WQI by Reach and Section (WY05-WY20)

<u>Annual</u>	LMV	UMV	MV	MG	LSB	USB	SB	LSDR	
<u>Avg.</u>	Reach	Reach	Section	Section	Reach	Reach	Section	Overall Avg.	
WY05	48	43	46	63	31	18	24	41/40	C (highest)
WY06	39	33	37	54	34	22	28	36/35	D+
WY07	36	28	33	49	40	27	34	36/35	D+
WY08	38	30	35	45	38	34	36	37/37	D+
WY09	38	29	34	45	38	32	35	36/36	D+
WY10	36	32	34	47	37	18	27	34/32	D
WY11	39	38	39	54	44	15	29	38/36	D+
WY12	35	35	35	47	39	9	24	33/31	D
WY13	37	32	35	44	35	11	23	32/30	D
WY14	18	19	18	36	28	11	19	22/20	E (lowest)
WY15	24	22	23	44	43	11	27	29/25	D
WY16	35	22	29	40	37	9	23	28/27	D
WY17	34	32	33	41	39	19	29	33/31	D
WY18	26	22	24	33	27	10	19	24/22	E+
WY19	36	30	34	42	35	14	24	31/30	D
WY20	37	33	36	44	41	15	28	34/32	D
<i>16-yr Norms</i>	<i>34.8</i>	<i>30.0</i>	<i>32.7</i>	<i>45.6</i>	<i>36.5</i>	<i>17.0</i>	<i>26.8</i>	<i>32.8/31.3</i>	<i>D Marginal</i>
<u>Winter</u>	LMV	UMV	MV	MG	LSB	USB	SB	LSDR Overall	
WY05	63	65	64	84	44	33	39	58/58	B (highest)
WY06	54	50	52	60	40	29	35	47/46	C
WY07	49	42	46	61	55	40	48	50/47	B-/C+
WY08	56	47	52	54	52	52	52	52/52	B
WY09	57	48	53	61	54	49	52	54/53	B
WY10	54	53	54	66	54	28	41	51/49	B-/C+
WY11	57	56	56	66	54	27	40	52/50	B-
WY12	48	49	49	58	44	14	29	43/41	C
WY13	58	53	56	67	49	21	35	50/48	B-/C+

WY14	26	26	26	55	39	15	27	32/29	D (lowest)
WY15	33	29	31	58	53	11	32	37/32	D+/D
WY16	44	38	41	57	52	14	33	41/37	C/D+
WY17	53	58	55	66	60	35	48	54/53	B
WY18	38	37	38	58	41	16	29	38/36	C/D+
WY19	58	56	57	69	58	29	43	54/52	B
WY20	54	58	55	63	54	19	37	50/47	B-/C+
<i>16-yr Norms</i>	<i>50.1</i>	<i>47.7</i>	<i>49.1</i>	<i>62.7</i>	<i>50.1</i>	<i>27.1</i>	<i>38.6</i>	<i>47.5/45.8</i>	<i>C+ Fair</i>
Summer	LMV	UMV	MV	MG	LSB	USB	SB	LSDR Overall	
WY05	31	24	28	45	20	5	13	25/24	D-/E+
WY06	23	14	19	44	30	18	24	26/23	D-/E+
WY07	23	14	19	34	24	14	19	22/20	E
WY08	23	20	22	31	25	18	21	23/22	E
WY09	21	14	18	31	25	16	20	21/20	E
WY10	21	17	20	33	26	9	17	21/19	E
WY11	23	17	20	37	30	5	17	22/20	E
WY12	22	18	20	25	27	4	15	19/17	E
WY13	18	14	16	18	23	5	14	16/14	E
WY14	10	11	10	12	16	9	12	11/11	F+
WY15	15	11	13	32	37	9	23	21/17	E
WY16	18	7	13	18	19	5	12	13/11	E-/F+
WY17	20	16	18	20	22	11	17	18/17	E
WY18	12	8	10	9	15	6	10	10/9	F (lowest)
WY19	23	11	18	23	22	3	13	16/14	E
WY20	25	17	21	29	29	10	20	22/21	E
<i>16-yr Norms</i>	<i>20.4</i>	<i>14.5</i>	<i>17.9</i>	<i>27.4</i>	<i>24.4</i>	<i>9.3</i>	<i>16.8</i>	<i>19.2/18.0</i>	<i>E Poor</i>

Table 5.1 WQI Letter/Color Code: A (>75) Very Good (dark blue), B (50-74) Good (light blue), C (38-49) Fair (green), D (25-37) Marginal (yellow), E (13-24) Poor (brown), and F (0-12) Very Poor (red). WQI values in red are below 16-yr norms (expressed in black italics) for the same reach or section of the river; values at 16-yr norms are in blue, values above in black. Overall LSDR WQI values are unweighted averages (each site value considered equal weight) and flow-weighted averages.

The running averages, as well as variances in monthly index values, for each reach of the lower river watershed are presented in the series of charts (5.1 through 5.6) on pages 19 and 20.

As shown on **Chart 5.1**, average annual WQI values associated with the **Lower Mission Valley Reach** (Sites 1-4) of the river have varied from a high of 48 (B+ Good) in WY05 to a low of 18 (E Poor) in WY14. The general trend in running average WQI for the reach, as well as for four individual monitoring sites, declined from the mid 40's (C Fair) during WY's '05 and '06 to the mid-teens (E Poor) by early WY15. The running average WQI (*black line*) improved to the mid-30's during the second half (April-Sept) of WY16

and climbed slightly higher during last year and WY20. Site 3 (Fashion Valley Mall, *blue line*) has consistently exhibited the lowest running average WQI, while Site 4 (FSDRIP at Mission Valley Rd., *red line*) has consistently witnessed the highest values for the reach. The most significant decline in the WQI for the reach over the 16-year monitoring period occurred in WY14. There was a steady, general improvement from WY14 lows during the second half of WY15 and throughout WY16 into WY17. A general decline occurred throughout WY18, followed by recovery to WY17 values in WY19 and WY20. The running average index for this reach has declined by approximately ten percent (from 45 to 35) over the 16-year monitoring period.

As shown in **Chart 5.2**, the range in monthly WQI values for the **Upper Mission Valley Reach** (Sites 5-7) of the river are similar to those in Lower Mission Valley, although somewhat less variable. Site 6 (Kaiser Ponds at Mission Valley Rd, *green line*) has continuously presented lowest running average WQI values since early 2017, while Site 7 (Admiral Baker Field at Zion, *blue line*), situated just upstream of the ponds, has presented the highest values on an extended basis since mid-2008. The highest average annual WQI reading of 65 (A Very Good) for the Upper Mission Valley reach was in WY05, whereas the lowest reading of 19 (E Poor) was in WY14. The overall trend in running average WQI values (*black line*) from mid 2010 through 2013 was generally positive. Index values for each site and for the entire reach that trended downward through WY18 have recovered to prior year levels in WY19 and WY20. The overall trend since WY06 has been negative (in decline) as growth of invasive aquatic plants and increase in biomass has proliferated throughout much of this reach during extended periods of minimal flow. The rate of decline in running average index in this reach over 16 years is about ten percent, decreasing from 40 in WY05 to the present value of 32. Significant recovery in this reach is problematic without improved channel maintenance due to extensive accrual of biomass and insufficient flushing.

Running average WQI for the **Mission Gorge Reach** (Sites 8-10) of the river, as shown in **Chart 5.3**, has also declined, especially during WY12- through WY14. Highest annual WQI values of 63 (A Very Good) occurred in WY05, contrasted with a low of 33 (C Marginal) in WY18. In general running average WQI for this reach is the highest of the five reaches with an average WQI of 45.6 (B Good). The trend in Mission Gorge WQI values (*black line*) are, however, comparable to those in the Mission Valley reaches. General decline in index values from WY06 through WY09, followed by a slight upturn in WY10 and WY11, and a more significant decline in subsequent water years to a low of 33 (D Marginal) in early WY15. WY17 witnessed an overall nine-point recovery in the running average WQI. The index for this reach fell during the second half of WY18 to a record low of 32. WY19 saw recovery to a high of 42 and to 44 by the end of WY20. The overall index has declined nearly 20 points (from 60 down to 40) over 16 years in this section of the river.

The **Lower Santee Basin Reach** (Sites 11,15 & 12) WQI values and running averages are shown in **Chart 5.4**. The range from winter month highs in the 50-70 range (B Good) to summer lows in the 10-15 range (E Poor) are common. Water quality improved in this reach from WY06 through WY11, then declined in subsequent water years, reaching a running average low of 27 (D-Marginal) in 2015, before recovering to the mid-40s (C Fair) throughout WY16 and low 40's in WY17. The previous low was surpassed by one point in both August and September of WY18. WY19 witnessed partial recovery to the mid 30's reaching 40 in WY20. Completion of the Forester Creek enhancement project (indicated by the *blue line*) extending from Prospect Ave. to the Mission Gorge Rd. has had a significant impact on overall river quality (*black line*) in the Lower Santee Basin portion of the river system. With above normal rainfall experienced in WY19 and WY20, the Lower Santee Basin running average index has improved to values comparable to those experienced in WY07 through WY11. The overall change in the index from WY05 through WY20 is less than one percent. This reach of the river has shown the least change in water quality metrics over the 16 years of monitoring, due in large part to Forester Creek improvements and permitted releases of reclaimed water from Santee Lakes.






Chart 5.5 on the previous page presents monthly and running average WQI values for the **Upper Santee Basin Reach** (Sites 13 & 14) of the river. This reach presents the poorest water quality values of all sections of the lower river system. Monthly values have seldom exceeded 20 (E Poor) since the summer of 2011 and are often less than 12 (F+ Very Poor) throughout all but wet-weather, winter months. The running average WQI for this reach has declined from highs above 30 (D Marginal) in WY09 to continuously between 10 and 12 (F Very Poor) during the five year period (WY12-WY16). WY17 saw a noticeable increase (ten points) in the running average index from early in the year reaching 18 (E- Poor) in September, however WY18 witnessed a reversal with a steady decline toward previous lows. WY19 witnessed partial recovery to prior highs, especially at site 14. The greatest variability has been associated with site 13, Mast Park East (*green line*). The reach index has fallen 44% (from 32 to 18) over the past decade presenting the greatest decline in running average WQI of all reaches. Advanced eutrophication of multiple ponds situated within and upstream of Mast Park has led to high levels of oxygen depletion throughout the year.

The monthly and running average variation in WQI values for the three main sections of the lower river (i.e., Santee Basin, Mission Gorge and Mission Valley) and the overall **Lower San Diego River** system (weighted average of all 15 monitoring sites) are presented in **Chart 5.6**. WQI running average values recovered from WY14 lows in all three sections of the river during WY15 through WY17. Values noticeably declined in WY18 then rebounded (to WY13 & WY16 levels) in WY19 and WY20 in all three sections of the lower river. The Mission Gorge section changed the least, while the upstream section (Santee Basin) the most. There were noticeable increases in index values in all three sections of the river and thus overall in WY20. The current LSDR running average WQI of 32 (D Marginal) is four percent above the 16-yr norm. The overall trend in running average WQI for the LSDR that remained relatively steady in the range of 35 to 40 between WY06 and WY12, but declined toward the low 20's in WY14 and early WY15, returned to the low 30's in WY16 and WY17 then dipped in WY18. The LSDR weighted running average index rose several points from 30 in WY19 to 32 over the past year. The overall running average decline has fallen approximately 10 points (from 40 to 30) over the 16 year period..

The overall decline in the index is a function of lowered oxygen levels in combination with elevated water temperatures and higher specific conductivities monitored at nearly all sites. These values are impacted by low streamflows especially during extended months without rainfall. WQI values are anticipated to measurably increase were streamflows to rise above current norms and effective aquatic growth abatement measures are implemented (or possibly occur through natural flushing) for specific reaches of the river. Higher minimum index values during the summer months result in positive gradients for 12-mo. running averages within a single water year, especially the case in the Mission Gorge section. Without interventions, overall negative trends in WQI values are expected to persist for many if not all portions of the lower river due to natural processes of eutrofication.

Depleted dissolved oxygen levels (often less than 3 mg/L) in conjunction with minimal dry-weather flow resulting in warmer water, higher SpC (more dissolved solids) are the primary causes of the low water quality index values. The low DO concentrations are believed to be the result of extensive and persistent eutrophication from buildup of organic-rich detritus combined with restricted water movement at a number of key monitoring sites. Until the spread of creeping water primrose (*Ludwigia grandiflora*)* and other invasive aquatics can be effectively curtailed and the effects of eutrophication better managed, water quality in several reaches of the lower river system is expected to remain significantly below that found in other reaches of the river where improved circulation, mixing and natural re-oxygenation occurs.

* *Ludwigia peploides*, *L. grandiflora*, *L. hexapetala* are members of a highly productive emergent aquatic perennial native to South and Central America, parts of the USA and likely Australia (USDA-ARS, 1997). It was introduced in France in 1830 and has become one of the most damaging invasive plants in that country. It has been more recently introduced to areas beyond its native range in the United States where it is often considered a noxious weed (INVADERS, 2009; Peconic Estuary Program, 2009). *L. grandiflora*, *et. al.* are adaptable and tolerate a wide variety of habitats where they can transform ecosystems both physically and chemically. It sometimes grows in nearly impenetrable mats; can displace native flora and interfere with flood control and drainage systems, clog waterways and impacts navigation and recreation. The plant also has allelopathic properties that can lead to dissolved oxygen crashes, the accumulation of sulphide and phosphate, 'dystrophic crises' and intoxicated ecosystems (Dandelot et al., 2005). Its common name is "floating water primrose", it produces a distinctive small yellow flower during its bloom cycle (May-Nov.). It is a perennial herb (a dicot) called marsh purslane; a member of family ORAGRACEAE.

California Invasive Plant Council (CALIPC) website

